

Kansuinine B Datasheet

4th Edition (Revised in July, 2016)

[Product Information]

Name: Kansuinine B

Catalog No.: CFN92866

Cas No.: 57685-46-8

Purity: >=95%

M.F: C₃₈H₄₂O₁₄

M.W: 722.7

Physical Description: Powder

Synonyms:

HO O H O OH

[Intended Use]

- 1. Reference standards;
- 2. Pharmacological research;
- 3. Synthetic precursor compounds;
- 4. Intermediates & Fine Chemicals;
- 5. Others.

[Source]

The roots of Euphorbia kansui.

[Biological Activity or Inhibitors]

Kansuinine A and B, from E. KANSUI have an inhibitory effect on IL-6-induced Stat3

activation by activating ERK1/2, inhibition of MEK significantly blocks the effects of them

on IL-6-induced Stat3 activation and tyrosine phosphorylation, suggests that blocking of

IL-6-induced signal transduction is partially due to the sustained activation of ERK1/2 by

kansuinine A and B, which in turn results in an increase of Stat3 serine phosphorylation

and SOCS-3 expression.[1]

Kansuinine A and B are toxic ingredients in Kansui Radix, the contents of them decreased

after being processed with vinegar .[2]

[Solvent]

Chloroform, Dichloromethane, Ethyl Acetate, DMSO, Acetone, etc.

[HPLC Method]^[2]

Mobile phase: Methanol- H2O ,gradient elution ;

Flow rate: 1.0 ml/min;

Column temperature: 38 °C;

The wave length of determination: 235 nm.

[Storage]

2-8℃, Protected from air and light, refrigerate or freeze.

[References]

[1] Chang J S, Lee S W, Park M H, et al. Planta Med., 2010, 76(14):1544-9.

[2] Li Z J, Li Y, Lan G, et al. Chinese Traditional Patent Medicine, 2011, 33(12):2122-5.

[Contact]

Address:

S5-3 Building, No. 111, Dongfeng Rd.,

Wuhan Economic and Technological Development Zone,

Wuhan, Hubei 430056,

Email: info@chemfaces.com

Tel: +86-27-84237783 Fax: +86-27-84254680

Web: www.chemfaces.com